

Dec 2015

- i) Defⁿ 1.24- A matrix E is in (row) echelon form if it has the following two properties:
- (i) The zero rows, if any, occur at the bottom
 - (ii) each leading (non zero) entry in a row is in a column to the right of the leading entry of the row above it
- ii) Defⁿ 1.25- An echelon matrix E is said to be in reduced echelon form if, in addition,
- (i) in each non-zero row, the leading entry is 1, and
 - (ii) in each col ~~not~~ containing the leading entry of a row, all other entries are zero

b) $A = \begin{pmatrix} -1 & -2 & -2 & -1 \\ 1 & 2 & 0 & 3 \\ 2 & 4 & 3 & 3 \\ 3 & 6 & 5 & 4 \end{pmatrix}$

$\xrightarrow{r_4 \rightarrow r_4 - r_3 + r_1} \begin{pmatrix} -1 & -2 & -2 & -1 \\ 1 & 2 & 0 & 3 \\ 2 & 4 & 3 & 3 \\ 0 & 0 & 0 & 0 \end{pmatrix}$

$\xrightarrow{r_2 \rightarrow r_2 - 2r_1} \begin{pmatrix} -1 & -2 & -2 & -1 \\ 0 & 0 & -2 & 2 \\ 2 & 4 & 3 & 3 \\ 0 & 0 & 0 & 0 \end{pmatrix}$

$\xrightarrow{r_3 \rightarrow r_3 + 2r_1} \begin{pmatrix} -1 & -2 & -2 & -1 \\ 0 & 0 & -2 & 2 \\ 0 & 0 & -1 & 1 \\ 0 & 0 & 0 & 0 \end{pmatrix}$

$\xrightarrow{r_2 \rightarrow r_2 - 2r_3} \begin{pmatrix} -1 & -2 & -2 & -1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & -1 & 1 \\ 0 & 0 & 0 & 0 \end{pmatrix}$

$\xrightarrow{r_1 \rightarrow -r_1} \begin{pmatrix} 1 & 2 & 2 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & -1 \\ 0 & 0 & 0 & 0 \end{pmatrix}$

$\xrightarrow{r_3 \leftrightarrow r_4} \begin{pmatrix} 1 & 2 & 0 & 3 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & -1 \\ 0 & 0 & 0 & 0 \end{pmatrix}$

c) A not invertible as A cannot be reduced to I_4 (rows not LI)

(~~✓~~ ~~det(A)~~)

d) no, we've shown only two are LI by applying row reduction (row reduc is like simultaneous eqns)
we can build r_2, r_3 from r_1 and r_3

e) row space of A has basis

$$\{(1, 2, 0, 3), (0, 0, 1, -1)\} = \{a_1, a_2\}$$

Span + LI by row reduc

f) $B = \begin{pmatrix} 1 & 2 & 4 & -1 \\ 2 & 4 & 6 & 0 \end{pmatrix} = \begin{pmatrix} r_1 \\ r_2 \end{pmatrix}$

~~SP(1, 2, 4, -1)~~

$$r_1 = a_1 + 4a_2 \Rightarrow$$

$$r_2 = 2a_1 + 6a_2$$

~~$a \in \text{row}(A)$
 $a = \lambda_1 a_1 + \lambda_2 a_2$~~

$\text{row}(B) \subseteq \text{row}(A)$
~~row(A) \subseteq row(B)~~
~~row(A) $\not\subseteq$ row(B)~~

r_1, r_2 LI as not a LC of each other
 $\Rightarrow \dim(B) = \dim(A)$
 $\Rightarrow \text{row}(A) = \text{row}(B)$ by Thm 3.45

$$C = I_4$$

$$\dim(C) = 4 \quad \dim(A) = 2$$

\Rightarrow can't possibly have the same row space

g) $\text{rank}(A) = \# \text{ non zero rows in RREF of } A$
 $= 2 \text{ by } \text{Q6}$

h) ~~W ⊆ V is a subspace~~ Defn 2.11

A subspace W of V is a non-empty subset of V which itself forms a vec space under the same ops

i) $E_1 = \text{all Echelon } 3 \times 3 \text{ in } \mathbb{R}$

$E_2 = \text{reduced ech } 3 \times 3 \text{ in } \mathbb{R}$

$$E_1, E_2 \subseteq M_{3 \times 3}(\mathbb{R})$$

E_1 a subspace of $M_{3 \times 3}(\mathbb{R})$

* $E_1 \neq \emptyset$ $I_3 \in E_1$

* $\begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} -1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix} \in E_1$

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix} + \begin{pmatrix} -1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix} = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 2 \\ 0 & 0 & 0 \end{pmatrix} \notin E_1, \text{ as non zero rows not at bottom}$$

no, not closed under +

E_2 no

$\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \in E_2$

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} + \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} = \begin{pmatrix} 2 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 2 \end{pmatrix} \notin E_2$$

leading entries = 2 \neq 1

no, not closed under +

2)a) V_a vs F a field

i) LI

Defn 3.13 - $v_1, \dots, v_n \in V$ are LI if the only soln to

$$\alpha_1 v_1 + \alpha_2 v_2 + \dots + \alpha_n v_n = 0 \quad \alpha_i \in F$$

$$\text{is } \alpha_1 = \alpha_2 = \dots = \alpha_n = 0$$

ii) Defn 3.13 - $\{v_1, \dots, v_n\} \subseteq V$ is a spanning set if for $v \in V$ $\forall v \in V \exists$

$$\alpha_1, \dots, \alpha_n \in F \text{ st}$$

$$v = \alpha_1 v_1 + \dots + \alpha_n v_n$$

iii) Defn 3.23 - A set $B = \{v_1, \dots, v_n\}$ is a basis for V if they are both a spanning set for V and LI

$$S_1 = \{(a, b, a+b, 1) \mid a, b \in \mathbb{R}\}$$

$$\text{no } (1, 1, 2, 1), (0, 0, 0, 1) \in S_1,$$

$$(1, 1, 2, 1) + (0, 0, 0, 1)$$

$$= (1, 1, 2, 2) \notin S_1$$

(or no zero, zero in \mathbb{R}^4 is $(0, 0, 0, 0)$)

$$ii) S_2 = \{(a, b, c, d) \in \mathbb{R}^4 : a+b = c+d\}$$

$$\begin{pmatrix} \text{(by} \\ \text{Subspace} \\ \text{test}) \end{pmatrix} = \{(a, b, c, a+b-c) \in \mathbb{R}^4\}$$

$$a=b=c=0 \Rightarrow 0 \in S_2 \Rightarrow S_2 \neq \emptyset$$

$$(a, b, c, a+b-c), (a', b', c', a'+b'-c') \in S_2,$$

$$(a, b, c, a+b-c) + (a', b', c', a'+b'-c')$$

$$= ((a+a'), (b+b'), (c+c'), (a+a') + (b+b') - (c+c'))$$

$$\in S_1$$

$$\begin{aligned} & \lambda(a, b, c, a+b-c) \\ &= (\lambda a, \lambda b, \lambda c, \lambda(a+b-c)) \\ &= (\lambda a, \lambda b, \lambda c, (\lambda a) + (\lambda b) - (\lambda c)) \end{aligned}$$

$\in S_1$, $\Rightarrow \checkmark$ by SS test

iii) $S_3 = \{(a, b, a+b, a^2) \mid a, b \in \mathbb{R}\}$

$\nexists (1, 1, 2, 1), (1, 1, 2, 1) \in S_3$

$$(1, 1, 2, 1) + (1, 1, 2, 1) = (2, 2, 4, 2)$$

$$a=2 \quad a^2=4 \neq 2$$

$\Rightarrow \notin S_3$

vi) $W = \text{span} \left\{ \underbrace{(1, 0, 1, 0)}_{r_1}, \underbrace{(2, 3, 2, 2)}_{r_2}, \underbrace{(1, 1, 2, 1)}_{r_3}, \underbrace{(0, -2, 1, -1)}_{r_4} \right\}$

i) these span the set by def'

\Rightarrow remains to check LI

\nexists \Rightarrow put in matrix and perform row ops to check LI

$$\left(\begin{array}{cccc} 1 & 0 & 1 & 0 \\ 2 & 3 & 2 & 2 \\ 1 & 1 & 2 & 1 \\ 0 & -2 & 1 & -1 \end{array} \right) \xrightarrow{\substack{r_2 \rightarrow r_2 - 2r_1 \\ r_3 \rightarrow r_3 - r_1}} \left(\begin{array}{cccc} 1 & 0 & 1 & 0 \\ 0 & 3 & 0 & 2 \\ 0 & 0 & 1 & 1 \\ 0 & -2 & 1 & -1 \end{array} \right)$$

$$\xrightarrow{r_4 \rightarrow r_4 + r_2 - r_3} \left(\begin{array}{cccc} 1 & 0 & 1 & 0 \\ 0 & 3 & 0 & 2 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 \end{array} \right)$$

\Rightarrow not LI as a zero

row

r_4 can be built from r_1, r_2, r_3

ii) $\dim W \leq 4$ as atmost 3 LI vec in span
 $\Rightarrow (\text{basis}) \leq 4$

\Rightarrow this cannot be a basis as these
are 4 LI vec

d) $W = \{(a, b, 0, 0) \mid a, b \in \mathbb{R}\}$

$$(0, 0, 0, 0) \in W \Rightarrow W \neq \emptyset$$

$$(a, b, 0, 0), (c, d, 0, 0) \in W$$

$$(a, b, 0, 0) + (c, d, 0, 0)$$

$$= ((a+c), (b+d), 0, 0) \in W$$

$$\lambda \in F$$

$$\lambda(a, b, 0, 0) = (\lambda a, \lambda b, 0, 0) \in W$$

\Rightarrow W a subspace by SS test

$$(a, b, 0, 0) = a(1, 0, 0, 0) + b(0, 1, 0, 0)$$

$$\Rightarrow \{(1, 0, 0, 0), (0, 1, 0, 0)\} = \mathcal{B}$$

Span W

clearly L(I (standard basis subset))

$$\Rightarrow \dim W = |\mathcal{B}| = 2$$

e) V vs over \mathbb{F}

$U, W \subseteq V$ subspaces

$$\begin{aligned} * \quad \underline{v} \in U, W \Rightarrow \underline{v} \in U \cap W \\ \Rightarrow U \cap W \neq \emptyset \end{aligned}$$

$$* \quad \underline{v}, \underline{w} \in U \cap W$$

$$\begin{aligned} \Rightarrow \underline{v}, \underline{w} \in U & \quad \underline{v}, \underline{w} \in W \\ \Rightarrow \underline{v} + \underline{w} \in U & \quad \underline{v} + \underline{w} \in W \quad \text{both ss} \end{aligned}$$

$$\Rightarrow \underline{v} + \underline{w} \in U \cap W$$

$$* \quad \underline{v} \in U \cap W \quad \lambda \in \mathbb{F}$$

$$\Rightarrow \underline{v} \in U \quad \underline{v} \in W$$

$$\Rightarrow \lambda \underline{v} \in U \quad \lambda \underline{v} \in W \quad \text{both ss}$$

$$\Rightarrow \lambda \underline{v} \in U \cap W$$

\Rightarrow Subspace by ss test

$$U = \{(a, 0, 0, 0) \mid a \in \mathbb{R}\}$$

$$W = \{(0, 0, 0, b) \mid b \in \mathbb{R}\} \quad \text{exclusive or}$$

$$U \cup W = \{(a, b, c, d) \mid b=c=0, a \neq 0 \text{ or } d \neq 0\}$$

$$(1, 0, 0, 0) \in U \quad (0, 0, 0, 1) \in W$$

$$\Rightarrow (1, 0, 0, 0), (0, 0, 0, 1) \in U \cup W$$

$$(1, 0, 0, 0) + (0, 0, 0, 1)$$

$$=(1, 0, 0, 1) \notin U \cup W$$

as $\notin U$ last coord $\neq 0$

$\notin W$ 1st $\neq 0$

3)a)

i) defn 4.1 - v, w vs on same field F $T: V \rightarrow W$ each $v \in V$ ass a vec $T(v) \in W$ is a L_T if

$$\text{i)} \quad T(v+v) = T(v) + T(v) \quad \forall v, v \in V$$

$$\text{ii)} \quad T(\alpha v) = \alpha T(v) \quad \forall v \in V \quad \forall \alpha \in F$$

ii) 4.2 - all vec v mapped to zero of w by T

$$\ker T = \{v \in V \mid T(v) = \underline{0}\}$$

iii) 4.18 - image of $T: V \rightarrow W$ i.e. 1consists of all images $T(v)$ of vec in V under T

$$\text{im } T = T(V) = \{T(v) \mid v \in V\}$$

b) $\ker T$

$$\star \quad T(\underline{0}) = \underline{0}$$

$$T(\underline{0} \underline{0}) = \underline{0} T(\underline{0}) = \underline{0}$$

 $\Rightarrow \underline{0} \in \ker T \Rightarrow \ker T \neq \emptyset$
* $\underline{0}, v \in \ker T$

$$\begin{aligned} T(v+v) &= T(v) + T(v) \\ &= \underline{0} + \underline{0} \\ &= \underline{0} \end{aligned} \quad \downarrow v, v \in \ker T$$

 $\Rightarrow v+v \in \ker T$
* $v \in \ker T \quad \lambda \in F$

$$T(\lambda v) = \lambda T(v) = L_T$$

$$= \lambda \underline{0}$$

$$= \underline{0}$$

 $\Rightarrow \lambda v \in \ker T$

\Rightarrow Subspace by SS test

C) i) $T: \mathbb{R}^4 \rightarrow \mathbb{R}^3$

$$T \begin{pmatrix} x \\ y \\ z \\ t \end{pmatrix} = \begin{pmatrix} x+y+z \\ x-t \\ y+z+t \end{pmatrix}$$

$$T \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} \quad \checkmark$$

$$T \begin{pmatrix} 0 \\ 1 \\ 0 \\ 0 \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix}$$

$$T \begin{pmatrix} 0 \\ 0 \\ 1 \\ 0 \end{pmatrix} = \begin{pmatrix} 0 \\ 1 \\ 1 \end{pmatrix}$$

$$T \begin{pmatrix} 0 \\ 0 \\ 0 \\ 1 \end{pmatrix} = \begin{pmatrix} 0 \\ -1 \\ 1 \end{pmatrix}$$

$$\text{Mat}(T) = \begin{pmatrix} 1 & 1 & 1 & 0 \\ 1 & 0 & 0 & -1 \\ 0 & 1 & 1 & 1 \end{pmatrix}$$

ii) Perform row ops to find basis
 for row space of $\text{Mat } T^T$
 $= \text{col space } \text{Mat } T$
 $= \text{im of } T$

$$\text{mat}(\tau)^T = \begin{pmatrix} 1 & 1 & 0 \\ 1 & 0 & 1 \\ 1 & 0 & 1 \\ 0 & -1 & 1 \end{pmatrix}$$

$$r_3 \rightarrow r_3 - r_2 \rightarrow \begin{pmatrix} 1 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & -1 & 1 \end{pmatrix}$$

$$r_2 \rightarrow r_2 - r_1 \rightarrow \begin{pmatrix} 1 & 1 & 0 \\ 0 & -1 & 1 \\ 0 & 0 & 0 \\ 0 & -1 & 1 \end{pmatrix}$$

$$r_4 \rightarrow r_4 - r_2 \rightarrow \begin{pmatrix} 1 & 1 & 0 \\ 0 & -1 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

$$r_2 \rightarrow -r_2 \rightarrow \begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & -1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

$$r_1 \rightarrow r_1 - r_2 \rightarrow \begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & -1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

$$\Rightarrow \text{Image} = \text{span} \left\{ \begin{pmatrix} 1 \\ 0 \\ 1 \\ -1 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \\ 0 \\ 0 \end{pmatrix} \right\}$$

(I + Spanning by row reduce
 \Rightarrow also a basis)

~~for~~

$$\begin{pmatrix} x \\ y \\ z \\ t \end{pmatrix} \text{ there is } \Leftrightarrow T \begin{pmatrix} x \\ y \\ z \\ t \end{pmatrix} = \underline{0}$$

$$\Leftrightarrow \begin{aligned} x + y + z &= 0 \\ x - t &= 0 \quad \Rightarrow x = t \\ y + z + t &= 0 \end{aligned}$$

$$y + z + t = 0$$

~~t ≠ 2x + y + z~~

$$z = -y - t = -y - x$$

$$\Rightarrow \begin{pmatrix} x \\ y \\ z \\ t \end{pmatrix} \text{ there is } \Leftrightarrow \begin{pmatrix} x \\ y \\ z \\ t \end{pmatrix} = \begin{pmatrix} x \\ y \\ -y-x \\ x \end{pmatrix}$$

$$\begin{pmatrix} x \\ y \\ -(y+x) \\ x \end{pmatrix} = x \begin{pmatrix} 1 \\ 0 \\ -1 \\ 1 \end{pmatrix} + y \begin{pmatrix} 0 \\ 1 \\ -1 \\ 0 \end{pmatrix}$$

$\Rightarrow \begin{pmatrix} 1 \\ 0 \\ -1 \\ 1 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \\ -1 \\ 0 \end{pmatrix}$ span the ker
clearly $L\mathbb{T} \Rightarrow$ a basis

a) let $T: V \rightarrow W$ be a ~~top~~ $L\mathbb{T}$

rank nullity;

$$\text{Rank}(T) + \text{Nullity}(T) = \dim(V)$$

$$e) A = \{(x, 0, z, 0) \mid x, z \in \mathbb{R}\}$$

$$i) F, g : \mathbb{R}^4 \rightarrow \mathbb{R}^4$$

$$F : \mathbb{R}^4 \rightarrow \mathbb{R}^4$$

$$F \begin{pmatrix} x \\ y \\ z \\ t \end{pmatrix} = \begin{pmatrix} x \\ 0 \\ z \\ 0 \end{pmatrix}$$

$$\begin{aligned} F \left(\begin{pmatrix} x \\ y \\ z \\ t \end{pmatrix} + \begin{pmatrix} a \\ b \\ c \\ d \end{pmatrix} \right) &= F \begin{pmatrix} x+a \\ y+b \\ z+c \\ t+d \end{pmatrix} \\ &= \begin{pmatrix} x+a \\ 0 \\ z+c \\ 0 \end{pmatrix} \\ &= \begin{pmatrix} x \\ 0 \\ z \\ 0 \end{pmatrix} + \begin{pmatrix} a \\ 0 \\ c \\ 0 \end{pmatrix} \\ &= F \begin{pmatrix} x \\ y \\ z \\ t \end{pmatrix} + F \begin{pmatrix} a \\ b \\ c \\ d \end{pmatrix} \end{aligned}$$

$$\begin{aligned} F \left(\lambda \begin{pmatrix} x \\ y \\ z \\ t \end{pmatrix} \right) &= F \begin{pmatrix} \lambda x \\ \lambda y \\ \lambda z \\ \lambda t \end{pmatrix} \\ &= \begin{pmatrix} \lambda x \\ 0 \\ \lambda z \\ 0 \end{pmatrix} \\ &= \lambda \begin{pmatrix} x \\ 0 \\ z \\ 0 \end{pmatrix} \\ &= \lambda F \begin{pmatrix} x \\ 0 \\ z \\ 0 \end{pmatrix} \end{aligned}$$

$\Rightarrow F$ linear

$$\text{im } F = \left\{ \begin{pmatrix} x \\ 0 \\ z \\ 0 \end{pmatrix} \mid x, z \in \mathbb{R} \right\}$$
$$= A$$

i) $g: \mathbb{R}^4 \rightarrow \mathbb{R}^4$

$$g \begin{pmatrix} x \\ y \\ z \\ t \end{pmatrix} = \begin{pmatrix} 0 \\ y \\ 0 \\ t \end{pmatrix}$$

explain on day;

check linear

4)a) defⁿ s. i

$A \in M_{n \times n}(F)$ $\forall F^n \quad V \neq \{0\} \quad \text{Evec if}$

$$Av = \lambda v$$

for $\lambda \in F$

λ = Eval of A

b)i)

$$\cancel{A = \begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 1 \end{pmatrix}}$$

$$\det(A - \lambda I) = \det \begin{pmatrix} 1-\lambda & 0 & 1 \\ 0 & 1-\lambda & 0 \\ 1 & 0 & 1-\lambda \end{pmatrix}$$
$$= (1-\lambda)[(1-\lambda)^2 - 0] + 0$$
$$- 1(0 - (1-\lambda))$$

$$= (1-\lambda)[(1-\lambda)^2 + 1]$$

$$= (1-\lambda)[1-2\lambda+\lambda^2 + 1]$$

$$= (1-\lambda)[2-2\lambda+\lambda^2]$$

$$\lambda = 1$$

$$\lambda^2 - 2\lambda + 2 = 0$$

$$\lambda = \frac{2 \pm \sqrt{2^2 - 4(2)(1)}}{2}$$

$$= \frac{2 \pm \sqrt{4-8}}{2}$$

$$= \frac{2 \pm \sqrt{-4}}{2} = 1 \pm i$$

$$b) i) \quad A = \begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

$$\begin{aligned} \det(A - \lambda I) &= \det \begin{pmatrix} 1-\lambda & 0 & 1 \\ 0 & 1-\lambda & 0 \\ 0 & 0 & 1-\lambda \end{pmatrix} \\ &= (1-\lambda) \left[(1-\lambda)^2 - 0 \right] - 0 \\ &\quad + 1 \left[0 - 1(1-\lambda) \right] \\ &= (1-\lambda)(1-\lambda)^2 - (1-\lambda) \\ &= (1-\lambda) \left[1 - 2\lambda + \lambda^2 - 1 \right] \\ &= (1-\lambda) (\lambda^2 - 2\lambda) \\ &= (1-\lambda) \lambda (\lambda - 2) \\ \Rightarrow \lambda &= 1, 0, 2 \end{aligned}$$

$$\underline{\lambda = 1}$$

$$A \underline{v} = \underline{v}$$

$$\Rightarrow (A - I) \underline{v} = \underline{0}$$

$$\Rightarrow \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix} \begin{pmatrix} a \\ b \\ c \end{pmatrix} = \underline{0}$$

$$\Rightarrow \begin{pmatrix} c \\ 0 \\ a \end{pmatrix} = \underline{0}$$

$$\begin{aligned} \Rightarrow c &= a = 0 \\ \Rightarrow \underline{v} &\in \left\{ \begin{pmatrix} 0 \\ b \\ 0 \end{pmatrix} \mid b \in \mathbb{R} \right\} \quad \text{ie } \begin{pmatrix} 0 \\ b \\ 0 \end{pmatrix} \end{aligned}$$

$\lambda = 0$

$$A\underline{v} = \underline{0}$$

$$\Rightarrow \begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 1 \end{pmatrix} \begin{pmatrix} a \\ b \\ c \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}$$

$$\Rightarrow \begin{pmatrix} a+c \\ b \\ a+c \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}$$

$$\Rightarrow b=0 \quad a=-c$$

$$\Rightarrow \underline{v} \in \left\{ \begin{pmatrix} a \\ 0 \\ -a \end{pmatrix} \mid a \in \mathbb{R} \right\} \quad \text{ie } \begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix}$$

$\lambda = 2$

$$A\underline{v} = 2\underline{v}$$

$$\Rightarrow (A - 2I)\underline{v} = \underline{0}$$

$$\begin{pmatrix} -1 & 0 & 1 \\ 0 & -1 & 0 \\ 1 & 0 & -1 \end{pmatrix} \begin{pmatrix} a \\ b \\ c \end{pmatrix} = \underline{0}$$

$$\Rightarrow \begin{pmatrix} -a+c \\ -b \\ a-c \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}$$

$$\Rightarrow a=c \quad b=0$$

$$\Rightarrow \underline{v} \in \left\{ \begin{pmatrix} a \\ 0 \\ a \end{pmatrix} \mid a \in \mathbb{R} \right\} \quad \text{ie } \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix}$$

ii) P, D

$$P = \begin{pmatrix} 0 & 1 & 1 \\ 1 & 0 & 0 \\ 0 & -1 & 1 \end{pmatrix} \quad D = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 2 \end{pmatrix}$$

c)) ? Eval of A invert

Assume for # $\lambda = 0$

$$\Rightarrow A\underline{v} = \lambda \underline{v}$$

$$\Rightarrow A\underline{v} = 0\underline{v} = \underline{0}$$

$$\Rightarrow A\underline{v} = \underline{0}$$

A invert $\Rightarrow A^{-1}$ exists

$$\Rightarrow A^{-1}A\underline{v} = A^{-1}\underline{0}$$

$$\Rightarrow I\underline{v} = \underline{0}$$

$$\Rightarrow \underline{v} = \underline{0}$$

$\Rightarrow \#$ by defⁿ $\underline{v} \neq \underline{0}$

ii) $A\underline{v} = \lambda \underline{v}$

A invert $\Rightarrow A^{-1}$ exists

$$\Rightarrow A^{-1}A\underline{v} = A^{-1}\lambda \underline{v}$$

$$\Rightarrow I\underline{v} = A^{-1}\lambda \underline{v}$$

$$\Rightarrow \underline{v} = \lambda A^{-1}\underline{v}$$

$$\Rightarrow \lambda^{-1}\underline{v} = A^{-1}\underline{v} \quad \text{allowable by (i)}$$

$$\Rightarrow A^{-1}\underline{v} = \lambda^{-1}\underline{v}$$

\Rightarrow yes!